TIERED EVALUATION OF LABORATORY DEVELOPED TESTS POLICY

History: NYSDOH Wadsworth Center's Clinical Laboratory Evaluation Program (CLEP) and its Clinical Laboratory Reference System's (CLRS) scientific staff adopted a three-tiered model for the risk-based review and approval of laboratory-developed tests (LDT) beginning November 14, 2016.

Update: Effective April 17, 2023, CLEP and CLRS revised the policy regarding Lifestyle tests. This revised policy applies to <u>all</u> laboratories applying for or holding a New York State clinical laboratory permit and all assays that require submission as indicated on the Test Approval webpage, which can be found at http://www.wadsworth.org/regulatory/clep/clinical-labs/obtain-permit/test-approval.

LDTs that were conditionally approved prior to the date of implementation of this revised policy will continue to hold conditional approval.

Except for Clinical Trial and Lifestyle assays, LDTs proposed by laboratories that do <u>not</u> currently hold a NYS clinical laboratory permit or the appropriate permit category for the testing proposed will continue to require review <u>and</u> approval before testing on NYS specimens may commence. Conditional approval is not available for such assays. Approval can only be granted when the laboratory has met all requirements and is issued a permit (initial) or permit amendment (new categories). Please refer to the Clinical Laboratory Evaluation Program Guide to Requirements and Services available on our website at http://www.wadsworth.org/regulatory/clep/clinical-labs.

With respect to Clinical Trial and Lifestyle LDTs, a NYS clinical laboratory permit must be issued before testing on NYS specimens may commence. To obtain a NYS clinical laboratory permit all requirements must be met, including successful participation in an onsite survey demonstrating compliance with all NYS statues, regulations, and standards.

Features of the tiered evaluation process

- 1. All LDTs must be submitted to CLEP following the appropriate test submission guidelines or relevant form currently posted on the CLEP Test Approval webpage, and include the elements described therein. This includes a **Risk Attestation Form** containing a succinct summary of the assay and responses to the questions listed on the Form; a description of each question is provided at the end of this policy document. The summary and responses to the questions will inform the assignment of the LDT to a risk classification by CLRS staff.
- 2. A classification assignment (high, moderate, low, clinical trial, or lifestyle) will be made by CLRS staff as part of the existing CLEP LDT validation review and approval process described on the CLEP Test Approval webpage. Laboratories are generally notified of the risk assignment within three weeks of the submission of a complete package. Incomplete, vague, or unclear responses on the Risk Attestation Form may result in delayed review.
- 3. Questions about completing the Risk Attestation form can be submitted to <u>CLEPVAL@health.ny.gov</u>. However, the department is unable to perform pre-submission risk determinations because each risk classification is predicated upon the information provided in the Risk Attestation Form as assessed by the Panel based on the flow diagram below.

Classifications

- <u>High risk</u>: LDTs assessed as <u>High risk</u> will <u>NOT</u> receive Conditional Approval and testing cannot be offered on specimens from NYS until the CLRS review has been completed and full approval has been granted. Review of LDTs assessed as <u>High risk</u> will be prioritized. Laboratories will be required to respond to CLRS reviews within 60 business days to avoid inactivation of the application but can apply for a one-time 60-day extension. Requests for extension should be made to CLEPVAL@health.ny.gov. High risk will apply to all LDT packages submitted by laboratories that are pending a permit or the appropriate permit category, unless the package qualifies for either a Clinical Trial or Lifestyle classification, though all other applicable permit requirements must be met. This includes laboratories engaged in a distributive testing model where either or both laboratories are pending permit or amendment.
- Moderate risk: LDTs assessed as Moderate risk will receive Conditional Approval if the laboratory holds a permit in the appropriate category of testing. Laboratories may offer the test when notified by CLEP of the moderate risk classification and conditional approval. However, the department reserves the right to withhold or withdraw conditional approval at its discretion. LDTs assessed as Moderate risk may not be reviewed further if they are well-established and are not a key determinant or have low impact. However, only those that are reviewed may receive full approval. Laboratories will be required to respond to CLRS reviews within 60 business days to avoid the rescinding of conditional approval. A one-time 60-day extension may be granted upon request. Requests must be made to CLEPVAL@health.ny.gov.
- <u>Low risk</u>: LDTs assessed as <u>Low</u> risk will receive full approval and will not be subject to review by
 CLRS staff, <u>provided the laboratory holds a permit in the appropriate category of testing</u>. Laboratories
 will be able to offer the test once notified by CLEP of the <u>Low risk</u> classification and approval.
 However, the department reserves the right to withhold approval and/or require CLRS review of the test
 at its discretion.
- Clinical Trial: LDTs assessed as being performed solely for Clinical Trial purposes will receive notification from CLEP recognizing that the proposed test is part of a clinical trial and may be offered to clients. The clinical trial for which the LDTs are performed must be approved by the National Institutes of Health (NIH) or another relevant independent Institutional Review Board (IRB). For additional information, please visit: https://clinicaltrials.gov/ct2/manage-recs/fdaaa. If the assay is used for both diagnostic and clinical trial purposes, then this classification does not apply. If the assay qualifies for the Clinical Trial classification initially but is then proposed for diagnostic use in the future, a new comprehensive method validation package must be submitted.
- <u>Lifestyle</u>: LDTs assessed as <u>Lifestyle</u> will receive notification from CLEP that the test may be offered to clients with the disclaimer that it has not been evaluated by the NYS Department of Health. Relevant documentation from the method validation submission will be shared with the Centers for Medicare and Medicaid Services (CMS) for a determination on whether the proposed testing is subject to Clinical Laboratory Improvement Amendments (CLIA) regulations. If not, CLEP will make a separate, independent assessment on the necessity for a NYS clinical laboratory permit. If the assay is subject to CLIA or NYS law requiring a clinical laboratory permit, CLEP/CLRS will make an independent assessment for the determination of the Lifestyle classification.

Category	Submission required?	Initial Approval	Review required	Review Priority
High	Yes	None	Yes	High ¹
Moderate	Yes	Conditional ^{2,3}	Yes	Medium
Low	Yes	Full ^{2,3}	No ⁴	
Clinical Trial	Yes	N/A ^{2,4}	No ⁴	
Lifestyle	Yes	N/A ^{2,4}	N/A ⁴	

¹Submissions for laboratories pending a permit or permit category are automatically assigned High Risk unless the package meets the conditions for Clinical Trial or Lifestyle.

Definitions of terms used in for High, Moderate, and Low classifications

Well-established: The laboratory has demonstrated the ability to consistently submit complete and organized applications that adequately prove competence for development of LDTs with the same or similar technology and where an appropriate validation protocol is followed on a consistent basis over time; **AND** the methodology:

- a. has been previously approved by FDA or NYS; or
- b. has, without significant modifications, resulted in a meaningful clinical impact as described in multiple peer-reviewed publications.

Key determinant: the test result provides critical or essential information to

- 1. diagnose, and/or
- 2. indicate a greater likelihood of developing a disease or condition, and/or
- 3. indicate eligibility for a specific treatment, and/or
- 4. provide prognostic information that influences patient management or treatment decisions, and/or
- 5. provides information on treatment adherence and/or drug use.

Impact: the likelihood that an inaccurate test result will negatively impact a patient's condition or lead to patient morbidity/mortality. An LDT will have a high impact if an analytically or clinically inaccurate result leads to erroneous diagnosis and/or prediction of an inappropriate treatment, thereby increasing the risk of significant harm or death.

²Provided the laboratory holds the appropriate permit category.

³The department reserves the right to withhold approval at its discretion.

⁴The department reserves the right to review all applications at its discretion.

LDT Classifications (see flow diagram below)

• High risk:

- O An LDT that uses a methodology that is <u>not well-established in the submitting lab</u> and provides critical or essential information (<u>key determinant</u>) about a serious or life-threatening disease, disorder, or condition, whether the reported result, if inaccurate, could be used to support an incorrect diagnosis and/or an inappropriate clinical treatment that is likely to increase the risk of significant harm or death (<u>high impact</u>), or
- An LDT that uses a methodology that is not <u>well-established in the submitting lab</u>, does not provide critical or essential information (<u>key determinant</u>) about a serious or life-threatening disease, disorder, or condition, but the reported result, if inaccurate, could be used to support an incorrect diagnosis and/or an inappropriate clinical treatment that is likely to increase the risk of significant harm or death (<u>high impact</u>).

Moderate risk:

- An LDT that uses a <u>well-established</u> methodology and provides critical or essential information (<u>key determinant</u>) about a serious or life-threatening disease, disorder, or condition, whether the reported result, if inaccurate, could be used to support an incorrect diagnosis and/or an inappropriate clinical treatment that is likely to increase the risk of significant harm or death (<u>high impact</u>), or
- An LDT that uses a <u>well-established</u> methodology, does not provide critical or essential information (<u>key determinant</u>) about a serious or life-threatening disease, disorder, or condition, but the reported result, if inaccurate, could be used to support an incorrect diagnosis and/or an inappropriate clinical treatment that is likely to increase the risk of significant harm or death (<u>high impact</u>), or
- An LDT that uses a methodology that is not <u>well-established in the submitting lab</u>, but is not considered a <u>key determinant</u>, and an inaccurate reported result is not likely to support an incorrect diagnosis and/or an inappropriate clinical treatment (low impact).

Low risk:

An LDT that uses a <u>well-established</u> methodology, does not provide critical or essential information (<u>key determinant</u>) about a serious or life-threatening disease, disorder, or condition, and an inaccurate reported result is not likely to support an incorrect diagnosis and/or an inappropriate clinical treatment (<u>low impact</u>). Packages that are submitted under an approved exemption may also be classified as Low Risk. Laboratories may apply for an exemption from full method validation submission by following the instructions available at https://www.wadsworth.org/regulatory/clep/clinical-labs/obtain-permit/test-approval.

Clinical Trial:


An LDT that is being performed as part of clinical trial approved by NIH or another relevant independent IRB on specimens for participant enrollment or management where the results are reported and are used for clinical decision making. Examples of testing performed for participant management include those that influence enrollment (exclusion or inclusion), safety, toxicity, or dosing.

Lifestyle:

Generally, an LDT is considered a lifestyle test when the intended use makes no clinical claims or direct reference to diseases or conditions; relates to the maintenance or adoption of a general state of health or healthy activity; or relates to the widely accepted role of a healthy lifestyle in the management of certain chronic diseases or conditions. A claim that the healthy lifestyle choice(s) may play an important role in health outcomes should be generally accepted.

A **Lifestyle** test would include, but is not limited to, a test to detect the best diet for a genotype, risk for wrinkles and/or freckles, the best fitness plan, aptitude for learning languages, snacking profile, food metabolism traits, etc. The following intended uses are examples of tests that would **not** be considered **Lifestyle** tests: genetic markers for risk of chronic disease, identification of carrier status of variants for diseases known to be caused by genetic mutation, identification of variants suspected to be associated with disease states or risk of developing a disease. Additional information regarding the LDT may be required so the Department can ascertain the intended use claims and risk of harm to customers. If the proposed assay includes a panel of biomarkers or tests where a portion of the biomarkers or tests are deemed clinical <u>or diagnostic</u>, the validation package is not eligible for the Lifestyle classification. <u>The laboratory may divide the panel such that the clinical or diagnostic</u> are offered separately.

NYSDOH Policy for Tiered Evaluation of Laboratory Developed Tests (LDT)



6

Revised 4/2023