

NEW YORK STATE DEPARTMENT OF HEALTH

OSWEGO COUNTY
INDOOR RADON ESTIMATES
PERCENT HOMES ABOVE 4 PCI/L
FEBRUARY 2007

Town or city	homes measured	living area*		basement**	
		best estimate (%)	range# (%)	best estimate (%)	range# (%)
Albion town	1	1.2	(0.1 - 3.6)	11	(5 - 19)
Amboy town	0	1.2	(0.1 - 3.3)	11	(5 - 18)
Boylston town	0	1.2	(0.2 - 3.5)	11	(5 - 18)
Constantia town	10	1.2	(0.3 - 4.4)	8	(3 - 16)
Fulton city	23	1.3	(0.6 - 2.8)	12	(8 - 16)
Granby town	5	1.3	(0.3 - 3.7)	11	(5 - 17)
Hannibal town	7	1.3	(0.2 - 3.5)	12	(7 - 20)
Hastings town	14	0.6	(0.1 - 1.8)	9	(5 - 14)
Mexico town	8	1.7	(0.5 - 4.0)	13	(8 - 19)
Minetto town	8	0.5	(0.2 - 1.9)	9	(4 - 15)
New Haven town	2	1.4	(0.4 - 4.3)	13	(7 - 21)
Orwell town	7	6	(3 - 10)	23	(16 - 37)
Oswego city	31	0.6	(0.2 - 1.3)	8	(5 - 11)
Oswego town	17	2	(1 - 4)	14	(10 - 18)
Palermo town	4	1.2	(0.2 - 3.4)	11	(6 - 17)
Parish town	3	0.9	(0.2 - 3.1)	9	(4 - 15)
Redfield town	1	0.9	(0.1 - 2.8)	10	(4 - 17)
Richland town	8	1.7	(0.6 - 4.2)	13	(8 - 19)
Sandy Creek town	4	1.9	(0.6 - 5.1)	15	(9 - 23)
Schroepfel town	14	1.7	(0.6 - 3.5)	14	(9 - 20)
Scriba town	9	1.6	(0.5 - 3.6)	13	(8 - 19)
Volney town	9	1.7	(0.6 - 3.6)	13	(8 - 18)
West Monroe town	9	0.2	(0.1 - 1.2)	6	(1 - 12)
Williamstown town	3	0.6	(0.2 - 2.2)	9	(2 - 15)
Oswego County	197	1.7		12	

*Long-term measurement, 90 days to one year.

**Short-term measurement, typically 2 to 7 days.

Uncertainty range of one sigma corresponds to a 68% confidence interval.